


Simultaneous Weak Events

a new temporal analysis project


Vinay Kashyap

Harvard-Smithsonian Center for Astrophysics


Chandra HETGS+ACIS-S grating dispersed events




AU Mic : ObsID 17 ; HETGS+ACIS-S



Flares on HD 189733 that seem to be tied to planetary phase




Procyon : is there any variability?




Limitations in current analyses

- Likelihoods are constructed with no regard to data order
 - ignoring auto-regression and ICA/SCA
 - fluctuations in consecutive bins, or groups of like fluctuations require human intervention via residual analysis




$$\chi^2=214.0$$




Limitations in current analyses

- Likelihoods are constructed with no regard to data order
 - ignoring auto-regression and ICA/SCA
 - fluctuations in consecutive bins, or groups of like fluctuations require human intervention via residual analysis
- Coincidences cannot be evaluated non-parametrically in multiple data streams
 - we are more likely to believe that something is real if a signal is seen simultaneously in independent data streams


AU Mic : ObsID 17 ; HETGS+ACIS-S




AU Mic : ObsID 17 ; HETGS+ACIS-S




AU Mic : ObsID 17 ; HETGS+ACIS-S




AU Mic : ObsID 17 ; HETGS+ACIS-S



AU Mic : ObsID 17 ; HETGS+ACIS-S



AU Mic : ObsID 17 ; HETGS+ACIS-S




Simulation

- Generate 500 draws from $\mathcal{N}(0,1)$
 - Find all fluctuations at $>1, 1.5, 2, 2.5, 3$ sigma
- Repeat 100 times
 - During each repetition, check how often a similar fluctuation is coincident with original fluctuation
 - Compute average frequency of coincidence
- Repeat 100 times
- Compare coincidence frequency with nominal probability of seeing coadded fluctuations of same sizes


Type I Error:
 Fraction of fluctuations that exceed $k \sigma$

k	1	1.5	2	2.5	3
$\mathcal{N}(\cdot)$	0.16	0.067	0.022	0.006	0.0013
$\langle \mathcal{N}(\cdot) \mathcal{N}(\cdot) \rangle$	0.025	0.0045	0.0005	0.0004	$8 \cdot 10^{-7}$
$\mathcal{N}(\cdot) + \mathcal{N}(\cdot)$	0.078	0.017	0.002	0.00015	10^{-5}
$\mathcal{N}(\cdot)^2$	0.025	0.0045	0.0005	0.0005	$4 \cdot 10^{-6}$

Procyon : is there any variability?



Procyon : is there any variability?



Not the first time someone has tried to figure this out.

Stetson & Welch 1993

Lehner et al. 2010

Stetson & Welch 1993, AJ 105, 1813

- variability index for two simultaneous streams
- first compute variance-weighted means
- then compute $\delta\chi$ for each stream
- compute variability index as sum of $\delta\chi^{(1)*} \delta\chi^{(2)}$

Lehner et al. 2010, PASP 122, 959

- Not the first time someone has tried to figure this out.
- Lehner et al. constructed a rank-ordered p-value statistic to find occultation events.
- p-value defined as $p(Z > z_j = -\ln((\prod_{i=1}^T r_{ij}) / N_p^T))$
- Optimized for occultation events with no large time-scale trends in the intensities
- Requires that statistical noise is not large

Type I is (relatively) easy; Type II is not.

Still can't deal with grouped fluctuations.

Want to detect weak events in streams dominated by background and statistical noise.