

## G120.1+01.4

## 1 Summary

- Common Name: Tycho's
- Distance: 2.4 kpc ( **Chevalier et al., 1980** )
- Center of X-ray emission (J2000): ( 00 25 19.9, 64 08 18.2 )
- X-ray size: 8.7' x8.6'
- Description:

### 1.1 Summary of Chandra Observations

Sequence	Obs ID	Instrument	Exposure <sub>uf</sub> (ks)	Exposure <sub>f</sub> (ks)	Date Observed	Aimpoint (J2000) ( $\alpha$ , $\delta$ )
500002	115	ACIS-235678	48.4	48.4	2000-09-20	( 00 25 27.6, 64 09 14.7 )

Exposure<sub>uf</sub> → Exposure time of un-filtered event file

Exposure<sub>f</sub> → Exposure time of filtered event file

- Images are incomplete ( ACIS-S3 covers most of the remnant while some eastern part is covered by ACIS-S2.)
- No background light-curve filtering.

### 1.2 Chandra Counts and Fluxes

Region	Energy Range (keV)	Signal (counts)	Rate (counts s <sup>-1</sup> )	F <sub>x</sub> <sup>abs</sup> (ergs cm <sup>-2</sup> s <sup>-1</sup> )	F <sub>x</sub> (ergs cm <sup>-2</sup> s <sup>-1</sup> )	L <sub>x</sub> (ergs s <sup>-1</sup> )
total?	0.3 - 10.0	4.691e+06	9.697e+01	5.12e-10	1.99e-09	1.37e+36
( 115 )	0.3 - 2.1	4.055e+06	8.382e+01	3.01e-10	1.76e-09	1.21e+36
	2.1 - 10.	6.425e+05	1.328e+01	2.12e-10	2.36e-10	1.62e+35

- N<sub>H</sub> = 0.63 (10<sup>22</sup> cm<sup>-2</sup>)
- Assumed distance: 2.4 kpc ( **Chevalier et al., 1980** )
- nH was derived by fitting the low energy spectrum with one thermal plasma model.
- Flux value given above is NOT from whole SNR.

### 1.3 Nearby Sources

Obs ID	Position (J2000)	Size	Net Count	Count rate	Note
115	( 00 24 29.9, 64 09 03.1 )	< 3.8"	48.4	1.00e-03	
	( 00 24 37.1, 64 10 59.4 )	< 2.7"	70.6	1.46e-03	
	( 00 24 40.7, 64 20 32.1 )	< 16.2"	414.0	8.56e-03	
	( 00 24 51.4, 64 17 25.3 )	< 8.7"	71.2	1.47e-03	
	( 00 24 52.5, 64 11 32.5 )	< 2.7"	14.7	3.04e-04	
	( 00 25 02.5, 64 13 51.4 )	< 3.3"	33.4	6.90e-04	
	( 00 25 58.0, 64 19 49.1 )	< 20.1"	39.0	8.06e-04	
	( 00 26 21.5, 64 12 11.4 )	< 12.6"	130.0	2.69e-03	

- (note) 1. This nearby source list is incomplete.  
 All the above sources are originally from the "src2.fits" file which is distributed with standard chandra processing.  
 Only sources with significant count rate and which are clear to visual inspection are included.
2. The size given above is the size of the region used in detecting that source.
3. For each source, background was subtracted from annular region around the source.

### 1.4 References

- Chevalier et al., 1980 ApJ, 235, 186 : Optical spectrum
- Reynoso et al., 1997 ApJ, 491, 816 : VLA


## 2 Fit Detail

- See spectrum page for used regions.

### 2.1 Total:

- soft energy region(0.3-1.6keV) was fitted with thermal plasma model
- Abundance of O, Ne, Mg, Fe were thawed and abundance of other O-like element were set to 0.


```
source=(xswabs * xsvapec)
reduced  $\chi^2 = 115.712$ 
nh = 0.6348 1022/cm2
```



### 2.2 Total:

- Same as above, showing all the energy range


```
source=(xswabs * xsvapec)
reduced  $\chi^2 = 2662.81$ 
nh = 0.6348 1022/cm2
```



### 2.3 Total:

- Addition to above model, several gaussian and a power model were added.

```
source=(xswabs * ((((((xsvapec + gauss1d) + gauss1d) +
gauss1d) + gauss1d) + gauss1d) + powlaw1d))
reduced  $\chi^2 = 98.8075$ 
nh = 0.6348 1022/cm2
```




### 3 Chandra Images : Band Images


- Left : raw image, binned by 1x1 pixel
- Right : gaussian smoothed version of above ( $\sigma = 2$  pixel)

#### 3.1 Wide Band Images


Total : 300-10000 eV



Soft Band : 300-2100 eV




**Hard Band : 2100-10000 eV**





**3.2 Band images used in true color image.**



**Red : 300-950 eV**



**Green : 950-2650 eV**




**Blue : 2650-7000 eV**




### 3.3 Misc.


: 1250-1410 eV




: 1580-2080 eV



**: 4080-6160 eV**



**: 6160-6700 eV**










## 4 Chandra Images : True Color

- Individual images are adaptively smoothed.
- Warning : the adaptive smoothing process sometimes produces artifacts.
- convolution method : fft
- kernel type : gauss
- significance ( min , max ) : ( 3 , 5 )

RED : 300-950 eV  
 GREEN : 950-2650 eV  
 BLUE : 2650-7000 eV







## 5 Chandra Spectrum







- Images show Regions used to extract spectra
- Regions with red strikes are excluded



### 5.1 ObsID 115

- Background was subtracted from the region around the SNR.

total?









## 6 Radio Image

- left : radio image
- right : chandra x-ray image with radio contour lines

### 1.375 GHz

- Image from **Reynoso et al.(1997)**
- 1 GHz flux density: 56 Jy (citepGREEN2001)




---



---

#### Summary of Observation

---

Telescope .....	VLA
Date .....	1994 Mar, Aug, Dec, 95 May
Frequency ....	1.375 GHz
Beam size ....	1.45x1.38"
1 sigma noise ..	0.14 mJy / beam

---




---


## 7 Images from Survey Missions

- Left : Chandra Image (0.3-10. keV)
- Center : Images from *SkyView* with the **same** scale
- right : Images from *SkyView* with a **reduced** scale


### ROSAT PSPC (1.0 deg): X-ray (0.1-2.4 keV)




### IRAS 12 micron: Infrared (12 micron)




### IRAS 25 micron: Infrared (25 micron)




**IRAS 60 micron: Infrared (60 micron)**




**IRAS 100 micron: Infrared (100 micron)**




**CO survey: Radio (115 GHz)**



**NRAO VLA Sky Survey (NVSS): Radio (1.4 GHz Continuum)**



**Westerbork Northern Sky Survey (WENSS): Radio (325 MHz Continuum)**



**Digitized Sky Survey: Optical (J or E band images with a few exceptions)**

